arXiv:math/0703041v1 [math.RT] 1 Mar 2007

INDECOMPOSABLE LINEAR GROUPS

V.A. Bovdi, V.P. Rudko

Dedicated to the memory of Professor A.V. Roiter

Abstract. Let G be a noncyclic group of order 4, and let $K = \mathbb{Z}$, $\mathbb{Z}_{(2)}$ and $\mathbb{Z}_2$ be the ring of rational integers, the localization of $\mathbb{Z}$ at the prime 2 and the ring of 2-adic integers, respectively. We describe, up to conjugacy, all of the indecomposable subgroups in the group $GL(m, K)$ which are isomorphic to G.

The first explicit description of the $\mathbb{Z}$-representations of the noncyclic group G of order 4 was obtained by L. Nazarova [4,5]. This description was significantly simplified and clarified after the discovery of the connection between the $\mathbb{Z}$-representations of the group G and the representations of a certain graph with five vertices defined on linear spaces over the field of two elements [2].

Let $K = \mathbb{Z}$, $\mathbb{Z}_{(2)}$, or $\mathbb{Z}_2$ be the ring of rational integers, the localization of $\mathbb{Z}$ at the prime 2 and the ring of 2-adic integers, respectively. Let $G = \langle a, b\rangle \cong C_2 \times C_2$ be a noncyclic group of order 4.

Let $\mathfrak{M}_n$ $(n > 1)$ be the set of all polynomials $f(x)$ of degree n over the field of two elements $\mathbb{F}_2$ which satisfy one of the following conditions:

- $f(x)$ is irreducible over $\mathbb{F}_2$;
- $f(x)$ is a power of a nonlinear irreducible polynomial over $\mathbb{F}_2$.

Define an action of the group

$$\begin{aligned} Aut(G) = \langle \sigma_1, \sigma_2 \mid &\sigma_1(a) = b, \quad \sigma_1(b) = a, \\ &\sigma_2(a) = a, \quad \sigma_2(b) = ab, \quad a, b \in G\rangle \cong S_3 \end{aligned}$$

on the set $\mathfrak{M}_n$ by $f^{\sigma_1}(x) = x^n f(x^{-1})$, $f^{\sigma_2}(x) = f(x+1)$, where $f(x) \in \mathfrak{M}_n$ and $n = deg(f(x))$.

By $\mathfrak{M}'_n$ we denote the set of the orbits of the set $\mathfrak{M}_n$ under the action of the group $Aut(G)$. Let

$$\begin{aligned} e_1 &= \tfrac{1}{4}(1+a)(1+b); \qquad e_2 = \tfrac{1}{4}(1-a)(1-b); \\ e_3 &= \tfrac{1}{4}(1+a)(1-b); \qquad e_4 = \tfrac{1}{4}(1-a)(1+b), \end{aligned} \tag{1}$$

1991 *Mathematics Subject Classification.* primary 20H15, secondary 20C10, 20C11.

The research was supported by OTKA No.T 037202 and No.T 038059

be the four primitive idempotents of the group algebra FG over a filed F containing K. The corresponding irreducible K-representations of these idempotents have the following form:

$$\begin{array}{lllllllllllll} \chi_1: & a & \mapsto & 1, & b & \mapsto & 1; & \chi_2: & a & \mapsto & -1, & b & \mapsto & -1; \\ \chi_3: & a & \mapsto & 1, & b & \mapsto & -1; & \chi_4: & a & \mapsto & -1, & b & \mapsto & 1. \end{array} \tag{2}$$

The representations χ_i $(i = 1, \dots, 4)$ are realized in the KG-module $KGe_i = Ke_i$ of the group algebra FG.

Consider the following K-representations Γ of the group $G = \langle a, b\rangle = C_2 \times C_2$ which were introduced in [3]. In the following we shell refer to them as $(\mathbb{L})$:

$$\Delta_n: \quad a \mapsto \begin{pmatrix} E_n & 0 & 0 & E_n & 0 \\ & 1 & 0 & 0 & 0 \\ & & -E_n & 0 & E_n \\ & & & -E_n & 0 \\ & & & & E_n \end{pmatrix}, \qquad b \mapsto \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ & E_n & 0 & 0 & E_n \\ & & -E_n & E_n & 0 \\ & & & E_n & 0 \\ & & & & -E_n \end{pmatrix};$$

$$W_0: \quad a \mapsto \begin{pmatrix} 1 & 1 & 0 & 1 \\ & -1 & 0 & 0 \\ & & 1 & 0 \\ & & & -1 \end{pmatrix}, \qquad b \mapsto \begin{pmatrix} 1 & 1 & 1 & 0 \\ & -1 & 0 & 0 \\ & & -1 & 0 \\ & & & 1 \end{pmatrix};$$

$$W_n: \quad a \mapsto \begin{pmatrix} D & 0 & 0 & 0 & 0 \\ & E_n & 0 & 0 & (\,0\ E_n\,) \\ & & -E_n & (\,0\ E_n\,) & 0 \\ & & & E_{n+1} & 0 \\ & & & & -E_{n+1} \end{pmatrix}, \qquad b \mapsto \begin{pmatrix} D & 0 & 0 & S & 0 \\ & E_n & 0 & (\,E_n\ 0\,) & 0 \\ & & -E_n & 0 & (\,E_n\ 0\,) \\ & & & -E_{n+1} & 0 \\ & & & & E_{n+1} \end{pmatrix};$$

$$T_n: \quad a \mapsto \begin{pmatrix} E_{n+1} & 0 & E_{n+1} & 0 \\ & -E_n & 0 & E_n \\ & & -E_{n+1} & 0 \\ & & & E_n \end{pmatrix}, \qquad b \mapsto \begin{pmatrix} 1 & & 0 & 0 & 0 \\ & E_n & 0 & 0 & E_n \\ & & -E_n & (\,E_n\ 0\,) & 0 \\ & & & E_{n+1} & 0 \\ & & & & -E_n \end{pmatrix};$$

$$\Delta_{n,1}: \quad a \mapsto \begin{pmatrix} E_n & 0 & (\,E_n\ 0\,) & 0 \\ & -E_{n+1} & 0 & E_{n+1} \\ & & -E_{n+1} & 0 \\ & & & E_{n+1} \end{pmatrix}, \qquad b \mapsto \begin{pmatrix} E_n & 0 & 0 & (\,0\ E_n\,) \\ & -E_{n+1} & E_{n+1} & 0 \\ & & E_{n+1} & 0 \\ & & & -E_{n+1} \end{pmatrix},$$

where $S = \left(\begin{smallmatrix} 0 & \dots & 0 & 1 & 1 \\ 0 & \dots & 0 & 0 & 0 \end{smallmatrix}\right)$, $\quad D = \left(\begin{smallmatrix} 1 & 1 \\ 0 & -1 \end{smallmatrix}\right)$.

The module M is called a module of the K-representation of the group G if M is a KG-module with finite KG basis.

Lemma 1. *If M is a module of the K-representation Γ of the group G and M does not contains regular direct summands, then*

$$(1 + a)(1 + b)M \subset 2M. \tag{3}$$

Proof. Let $\overline{K} = K/2K$ be the field of 2 elements. From the description of the $\overline{K}$-representations of group $G \cong C_2 \times C_2$ (see [1]) follows that if $\overline{M}$ is a module of indecomposable $\overline{K}$-representation of G and $\overline{M} \neq \overline{K}G$, then

$$(1+a)(1+b)\overline{M} = 0. \tag{4}$$

Let M be a module of an indecomposable K-representation of G and $M \neq KG$. The description of the K-representations of G (see [4,5]) shows that $\overline{M} = M/2M$ is a sum of indecomposable $\overline{K}G$-modules, and it is different from $\overline{K}G$. Hence from (4) follows (3), so the proof is complete.

Let M be a module of the K-representation of the group $G \cong C_2 \times C_2$, which satisfies the condition of Lemma 1. Put $M' = e_1M + e_2M + e_3M + e_4M \subset FM$, where $V_0 = M'/M$, $M_i = e_iM + M$ and $V_i = M_i/M$ $(i = 1, 2, 3, 4)$. Clearly, by Lemma 1 we get that $2M' \subset M$, $2M_i \subset M$ and V_0, V_i $(i = 1, 2, 3, 4)$ are vector spaces over the field $\mathbb{F}_2 = K/2K$. These spaces form a diagram

$$\begin{tikzcd}
V_1 \arrow[dr] & & V_3 \arrow[dl] \\
& V_0 & \\
V_2 \arrow[ur] & & V_4 \arrow[ul]
\end{tikzcd}$$

in which the arrows correspond to homomorphisms generated by the embeddings of M_i to M' $(i = 1, \dots, 4)$. Thus, the 5-tuple of linear spaces $(V_0; V_1, V_2, V_3, V_4)$ is a representation of the following graph on five points:



(5)

The following quadratic form

$$B(x) = x_0^2 + x_1^2 + x_2^2 + x_3^2 + x_4^2 - x_0(x_1 + x_2 + x_3 + x_4)$$

is defined on 5-dimension vectors corresponding to the graph (5), where $x = (x_0; x_1, x_2, x_3, x_4) \in \mathbb{Z}^5$.

A vector $x = (x_0; x_1, x_2, x_3, x_4) \in \mathbb{Z}^5$ is called a *positive root* of the form B if $B(x) = 1$ and $x_i \geq 0$, where $i = 0, \dots, 4$. The *dimension vector* $d(\Gamma) = (d_0; d_1, d_2, d_3, d_4)$ of a K-representation Γ of the group G is denoted by $d_i = dim_{\mathbb{F}_2} V_i$, where $V(\Gamma) = (V_0; V_1, V_2, V_3, V_4)$ is the 5-tuple of linear spaces corresponding to the representation Γ.

We know (see [2]), that if Γ is an indecomposable K-representation of the group G, which is different from the regular and irreducible representations, then either $B(d(\Gamma)) = 0$ or $d(\Gamma)$ is a positive root of the form $B(x)$ (i.e. $B(d(\Gamma)) = 1$). In addition, $d_0 > 0$ and $d_1 + d_2 + d_3 + d_4 > 1$ for any positive root $d = (d_0; d_1, d_2, d_3, d_4)$.

Moreover, for every positive root $d = (d_0; d_1, d_2, d_3, d_4)$ such that $d_0 > 1$ and $d_1 + d_2 + d_3 + d_4 > 1$ there exists only one indecomposable K-representation of the group G, such that the degree Γ is equal to $d_1 + d_2 + d_3 + d_4$.

Clearly, the one parameter family of indecomposable K-representations of the group G of degree $4m$ corresponds to the vector $d = (2n; n, n, n, n)$ (of course, $B(d) = 0$). The positive roots of the form B are listed in the following table:

$m = d_1 + d_2 + d_3 + d_4$	$d = (d_0; d_1, d_2, d_3, d_4)$	$t(m)$
$4n$ $n \geq 1$	$(2n + 1; n, n, n, n)$	1
$4n$ $n \geq 1$	$(2n - 1; n, n, n, n)$	1
$4n + 2$ $n \geq 1$	$(2n + 1; n + 1, n + 1, n, n)$	6
$4n + 1$ $n \geq 1$	$(2n; n + 1, n, n, n)$	4
$4n + 2$ $n \geq 1$	$(2n + 1; n + 1, n, n, n)$	4
$4n + 3$ $n \geq 1$	$(2n + 2; n + 1, n + 1, n + 1, n)$	4
$4n + 3$ $n \geq 1$	$(2n + 1; n + 1, n + 1, n + 1, n)$	4

Table 1.

where $t(m)$ denotes the number of roots with prescribed m and d_0. These roots are obtained from the root d by permutations of its components d_1, d_2, d_3, d_4.

First we describe the indecomposable K-representations Γ of G with common dimension vector $d(\Gamma) = (2n; n, n, n, n)$ (we recall that $B(d(\Gamma)) = 0$).

We will consider each polynomial $f(x) \in \mathbb{F}_2[x]$ as a polynomial over K, replacing $0, 1 \in \mathbb{F}_2$ by $0, 1 \in K$, respectively. Define $\widetilde{f(x)} = \begin{pmatrix} 0 & & & \alpha_0 \\ 1 & & & \alpha_1 \\ & \ddots & & \vdots \\ 0 & & 1 & \alpha_{n-1} \end{pmatrix}$ with the help of the corresponding matrix of $f(x) = x^n - \alpha_{n-1}x^{n-1} - \cdots - \alpha_1 x - \alpha_0 \in \mathbb{F}_2[x]$.

Each polynomial $f(x) \in \mathfrak{M}_n \cup \{x^n, (x+1)^n\}$, where $\mathfrak{M}_1 = \varnothing$, defines the following indecomposable K-representation of the group G:

$$\Delta_f : a \mapsto \begin{pmatrix} E_n & 0 & U_{11} & 0 \\ & -E_n & 0 & U_{22} \\ & & -E_n & 0 \\ & & & E_n \end{pmatrix}; \qquad b \mapsto \begin{pmatrix} E_n & 0 & 0 & U_{12} \\ & -E_n & U_{21} & 0 \\ & & E_n & 0 \\ & & & -E_n \end{pmatrix},$$

where only one of the matrices U_{ij} $(1 \leq i, j \leq 2)$ is equal to $\widetilde{f(x)}$, and the three other U_{ij} are equal to E_n.

Suppose that $\mathfrak{F}(f(x))$ is exactly that representation Δ_f for which $U_{12} = \widetilde{f(x)}$. For every K-representation Γ of the group G and for an automorphism $\phi \in Aut(G)$

we define a K-representation Γ^{ϕ} of the group G by $\Gamma^{\phi}(g) = \Gamma(\phi(g))$, where $g \in G$. The K-representations Γ and Γ^{ϕ} are called *conjugate.*

Lemma 2. *Let Γ be either an indecomposable K-representation from the list $(\mathbb{L})$ or its tensor product with the character χ_2 from (2). Then we have:*

Γ	$d(\Gamma) = (d_0; d_1, d_2, d_3, d_4)$
$\Delta_{n,1}$	$(2n+1; n, n+1, n+1, n+1)$
Δ_n^*	$(2n+1; n+1, n, n, n)$
Δ_n	$(2n+1; n+1, n, n, n)$
W_n	$(2n+1; n+1, n+1, n+1, n+1)$
T_n	$(2n+1; n+1, n, n, n+1)$
$\Delta_{n,1}^*$	$(2n+2; n, n+1, n+1, n+1)$
W_n^*	$(2n+3; n+1, n+1, n+1, n+1)$
$\Delta_n \otimes \chi_2$	$(2n; n, n+1, n, n)$
$(\Delta_n \otimes \chi_2)^*$	$(2n; n, n+1, n, n)$
$\Delta_{n,1} \otimes \chi_2$	$(2n+1; n+1, n, n+1, n+1)$
$(\Delta_{n,1} \otimes \chi_2)^*$	$(2n+2; n+1, n, n+1, n+1)$
$T_n \otimes \chi_2$	$(2n+1; n, n+1, n, n+1)$

Table 2.

Proof. We prove it for $\Gamma = \Delta_{n,1}$. Let

$$\{v_1, \dots, v_n, u_1, \dots, u_{n+1}, w_1, \dots, w_{n+1}, f_1, \dots, f_{n+1}\}$$

be a basis of the module M of the K-representation $\Delta_{n,1}$. Clearly,

$$e_1 v_i = \tfrac{1}{4}(1+a)(1+b)v_i = v_i; \qquad e_1 u_i = 0;$$
$$e_1 w_i = \tfrac{1}{4}(1+a)(2w_i + u_i) = \begin{cases} \frac{1}{2}v_i, & i \neq n+1; \\ 0, & i = n+1; \end{cases}$$
$$e_1 f_i = \tfrac{1}{4}(1+b)(2f_i + u_i) = \begin{cases} \frac{1}{2}v_{i-1}, & i = 2, \dots, n+1; \\ 0, & i = 1; \end{cases}$$
$$e_2 v_i = \tfrac{1}{4}(1-a)(1-b)v_i = 0; \qquad e_2 u_i = u_i;$$
$$e_2 w_i = \tfrac{1}{4}(1-a)u_i = \tfrac{1}{2}u_i; \qquad e_2 f_i = \tfrac{1}{4}(1-b)(-u_i) = -\tfrac{1}{2}u_i;$$

$$e_3 v_i = \tfrac{1}{4}(1+a)(1-b)v_i = 0; \qquad e_3 u_i = 0; \quad e_3 w_i = -\tfrac{1}{4}(1+a)u_i = 0;$$
$$e_3 f_i = \tfrac{1}{4}(1-b)(2f_i + u_i) = \begin{cases} f_i + \frac{1}{2}(u_i - v_{i-1}), & i = 2, \dots, n+1; \\ f_1 + \frac{1}{2}u_1, & i = 1; \end{cases}$$

$$e_4 v_i = \tfrac{1}{4}(1-a)(1+b)v_i = 0; \qquad e_4 u_i = 0;$$
$$e_4 w_i = -\tfrac{1}{4}(1-b)u_i = -\tfrac{1}{2}u_i; \quad e_4 f_i = -\tfrac{1}{4}(1+b)u_i = 0.$$

Since $V_i = (e_iM + M)/M$ we have that

$$\begin{aligned}
V_1 =& \langle \tfrac{1}{2}v_i + M \mid i = 1, \dots, n\rangle; \quad V_2 = \langle \tfrac{1}{2}u_i + M \mid i = 1, \dots, n+1\rangle;\\
V_3 =& \langle \tfrac{1}{2}u_1 + M, \quad \tfrac{1}{2}(u_i - v_{i-1}) + M \mid i = 2, \dots, n+1\rangle;\\
V_4 =& \langle -\tfrac{1}{2}u_i + M \mid i = 1, \dots, n+1\rangle.
\end{aligned}$$

Finally, it is easy to check that

$$V_0 = \langle \tfrac{1}{2}u_i + M,\ \tfrac{1}{2}v_j + M \mid i = 1, \dots, n+1;\ j = 1, \dots, n\rangle$$

and $d(\Delta_{n,1}) = (2n+1; n, n+1, n+1, n+1)$, so the proof is complete. The proofs for other representations Γ are similar to this one. The lemma is proved.

Lemma 3. *Let $K \in \{\mathbb{Z}, \mathbb{Z}_{(2)}, \mathbb{Z}_2\}$. Up to conjugacy, all indecomposable K-representations Γ of $G \cong C_2 \times C_2$, with common dimension vector $d(\Gamma) = (2n; n, n, n, n)$ belong to the either $\mathfrak{F}(f(x))$ $(f(x) \in \mathfrak{M}_n \cup \{x^n, (x+1)^n\})$ or to $\mathfrak{F}'(f(x)) = \mathfrak{F}(x^n) \otimes \chi_2$, where χ_2 is from (2).*

Up to equivalence, all indecomposable K-representations Γ of G with $d(\Gamma) = (2n; n, n, n, n)$ belong to one of the following types:

- $\mathfrak{F}(f(x))$, $\quad f(x) \in \mathfrak{M}_n \cup \{x^n, (x+1)^n\}$;
- $(\mathfrak{F}(x^n))^{\sigma_1}$;
- $\mathfrak{F}'(x^n)$;
- $(\mathfrak{F}'(x^n))^{\sigma_1}$;
- $(\mathfrak{F}'(x^n))^{\sigma_2}$, *where* $\sigma_1, \sigma_2 \in Aut(G)$.

Proof. The analysis of well known Nazarova's results on the representations of the group G (see [3,4]) shows that an indecomposable K-representation Γ of G, such that $d(\Gamma) = (2n; n, n, n, n)$, is equivalent to one of the representations Δ_f, where $f(x) \in \mathfrak{M}_n \cup \{x^n, (x+1)^n\}$, except the K-representation $(\mathfrak{F}'(x))^{\sigma_2}$, which is equivalent to the following representation:

$$a \mapsto \begin{pmatrix}
L_1 & 0 & 0 & 0 & 0 & 0\\
 & E_{n-1} & 0 & 0 & E_{n-1} & 0\\
 & & E_{n-1} & E_{n-1} & 0 & 0\\
 & & & -E_{n-1} & 0 & 0\\
 & & & & E_{n-1} & 0\\
 & & & & & -L_1
\end{pmatrix}; \quad b \mapsto \begin{pmatrix}
L_1 & 0 & 0 & & L_2 & 0\\
 & E_{n-1} & 0 & 0 & J_{n-1}(1) & L_3\\
 & & -E_{n-1} & -E_{n-1} & 0 & 0\\
 & & & E_{n-1} & 0 & 0\\
 & & & & E_{n-1} & 0\\
 & & & & & L_1
\end{pmatrix},$$

where $J_k(1)$ is a Jordan block of dimension k with ones in the main diagonal, $L_1 = \left(\begin{smallmatrix} 1 & 1\\ 0 & -1\end{smallmatrix}\right)$, $\quad L_2 = \left(\begin{smallmatrix} 1 & \dots & 0\\ 0 & \dots & 0\end{smallmatrix}\right)$, $\quad L_3 = \begin{pmatrix} 0 & 0\\ \vdots & \vdots\\ 0 & 1\end{pmatrix}$. So the lemma is proved.

In the next Theorem among other K-representations of the group G we list those indecomposable K-representations Γ such that $d(\Gamma)$ is a root of the form $B(x)$ (i.e. $B(d(\Gamma)) = 1$). Analyzing Nazarova's classification [4,5], we can see that any such obtained K-representation Γ can be derived from Ω and χ_i, (where Ω is from ($\mathbb{L}$), χ_i is from (2)) by one of the following operations:

- taking its conjugate Ω^{ϕ}, where $\phi \in Aut(G)$;
- taking the contragradient K-representation Ω^* $(\Omega^*(g) = \Omega^T(g^{-1}),\ g \in G)$;
- taking the tensor product $\Omega \otimes \chi_i$.

Our main result reads as follows:

Theorem. *Let $K = \mathbb{Z}$, $\mathbb{Z}_{(2)}$, or $\mathbb{Z}_2$ be the ring of rational integers, the localization of $\mathbb{Z}$ at the prime 2 and the ring of 2-adic integers, respectively. All indecomposable subgroups in the group $GL(m, K)$ which are isomorphic to the group $G = \langle a, b\rangle \cong C_2 \times C_2$ can be described, up to conjugacy, by groups which are generated by matrices $\Gamma(a)$ and $\Gamma(b)$, where Γ is a K-representation of G from the following table:*

m	Γ	$d(\Gamma)$	$St(\Gamma)$	$S(\Gamma)$
$4n+1$ $n \geq 1$	Δ_n	$(2n+1; n+1, n, n, n)$	$Aut(G)$	1
$4n+1$ $n \geq 1$	Δ_n^*	$(2n+1; n+1, n, n, n)$	$Aut(G)$	1
$4n+1$ $n \geq 1$	$\Delta_n \otimes \chi_2$	$(2n+1; n, n+1, n, n)$	$\langle\sigma_1\rangle$	3
$4n+1$ $n \geq 1$	$\Delta_n^* \otimes \chi_2$	$(2n; n, n+1, n, n)$	$\langle\sigma_1\rangle$	3
$4n+3$ $n \geq 0$	$\Delta_{n,1}$	$(2n+1; n, n+1, n+1, n+1)$	$Aut(G)$	1
$4n+3$ $n \geq 0$	$\Delta_{n,1}^*$	$(2n+2; n, n+1, n+1, n+1)$	$Aut(G)$	1
$4n+3$ $n \geq 0$	$\Delta_{n,1} \otimes \chi_2$	$(2n+1; n+1, n, n+1, n+1)$	$\langle\sigma_1\rangle$	3
$4n+3$ $n \geq 0$	$\Delta_{n,1}^* \otimes \chi_2$	$(2n+2; n+1, n, n+1, n+1)$	$\langle\sigma_1\rangle$	3
$4n+2$ $n \geq 1$	T_n	$(2n+1; n+1, n, n+1, n)$	$\langle\sigma_1\sigma_2\sigma_1\rangle$	3
$4n+2$ $n \geq 1$	$T_n \otimes \chi_2$	$(2n+1; n, n+1, n, n+1)$	$\langle\sigma_1\sigma_2\sigma_1\rangle$	3
$4n+4$ $n \geq 0$	W_n	$(2n+1; n+1, n+1, n+1, n+1)$	$Aut(G)$	1
$4n+4$ $n \geq 0$	W_n^*	$(2n+3; n+1, n+1, n+1, n+1)$	$Aut(G)$	1
$4n$ $n \geq 1$	$\mathfrak{F}(f(x))$, $(f(x) \in \mathfrak{M}'_n)$		$St[f(x)]$	$\frac{6}{\lvert St[f(x)]\rvert}$
$4n$ $n \geq 1$	$\mathfrak{F}(x^n)$		$\langle\sigma_2\sigma_1\sigma_2\rangle$	3
$4n$ $n \geq 1$	$\mathfrak{F}'(x^n) =$ $\mathfrak{F}(x^n) \otimes \chi_2$		$\langle\sigma_2\sigma_1\sigma_2\rangle$	3
4	Γ is the regular rep.			1

Table 3

Here $St(\Gamma) = \{\phi \in Aut(G) \mid \Gamma^{\phi}$ *is equivalent to* $\Gamma\}$ *is the stabilizer of* Γ *in* $Aut(G)$, $S(\Gamma) = \frac{|Aut(G)|}{|St(\Gamma)|}$ *is the number of non-equivalent* K*-representations which are conjugate to* Γ, Γ^* *is the contragradient* K*-representation of* Γ*, defined by* $\Gamma^*(g) = \Gamma^T(g^{-1}) \quad (g \in G)$.

Proof of the Theorem. Let G_1 and G_2 be subgroups in $GL(m, K)$ isomorphic to $G = \langle a, b\rangle \cong C_2 \times C_2$ with isomorphisms $\Gamma_i : G \mapsto G_i$, $(i = 1, 2)$. Then Γ_i is a faithful K-representation of G. By definition, the K-representation Γ_1 is conjugate to the K-representation Γ_2, if there are exist $C \in GL(m, K)$, $\phi \in Aut(G)$, such that $C^{-1}\Gamma_1(\phi(g))C = \Gamma_2(g)$, where $g \in G$. In other words, K-representations Γ_1 and Γ_2 of the group G are conjugate, if the groups $\Gamma_1(G)$ and $\Gamma_2(G)$ are conjugate in $GL(m, K)$. This implies the "if" part of the Theorem.

Now suppose G_1 and G_2 are conjugate in $GL(m, K)$ and $\tau : G_1 \mapsto G_2$ is an isomorphism such that $\tau(g) = C^{-1}gC$ for some matrix $C \in GL(m, K)$, where $g \in G$. Obviously, the map

$$\phi : G \xrightarrow{\Gamma_1} G_1 \xrightarrow{\tau} G_2 \xrightarrow{\Gamma_2^{-1}} G$$

is an automorphism of G and

$$\Gamma_2^{\phi}(g) = \Gamma_2(\Gamma_2^{-1}\tau\Gamma_1(g)) = \tau(\Gamma_1(g)) = C^{-1}\Gamma_1(g)C,$$

where $g \in G$, i.e. the K-representations Γ_1 and Γ_2 are conjugate.

Let Γ be a faithful K-representation of G, let $St(\Gamma)$ be the stabilizer of Γ in $Aut(G)$ and let $\{g_1, \dots, g_t\}$ be a system of representatives of the cosets of $Aut(G)$ by $St(\Gamma)$. Then $\{\Gamma^{g_i} \mid i = 1, \dots, t\}$ is the set of the pairwise nonequivalent K-representations of G, which are conjugated with Γ.

If Γ is an indecomposable K-representation of G and $d(\Gamma)$ is a root of the form $B(x)$, then $d(\Gamma^{\phi})$ $(\phi \in Aut(G))$ is also a root of the form $B(x)$. The vector $d(\Gamma^{\phi})$ can be obtained from the vector $d(\Gamma) = (d_0; d_1, d_2, d_3, d_4)$ by a permutation of the last four components of $d(\Gamma)$. It follows, that $S(\Gamma)$ (see Table 1) is the number of roots, which we obtained from $d(\Gamma)$ by permutation of some components in $d(\Gamma)$.

To finish the proof, we must compare the number of K-representations of degree m in Table 1 to the sum of all $t(m)$ for a given m in the Table 3, and then we can apply Lemma 3. So the lemma is proved.

Remark. *Let us substitute each* K*-representation* Γ *of* G *in Table 2 by* Γ^{σ}*. Here* σ *denotes a coset of* $Aut(G)$ *by* $St(\Gamma)$*. In this manner we obtain all faithful indecomposable* K*-representations of* G *up to equivalence.*

The proof of this *Remark* follows from the analysis of the classification of the representations described by L. Nazarova [3,4].

References

1. Bashev, V. A., *Representations of the group* $Z_2 \times Z_2$ *in a field of characteristic* 2*. (Russian)*, Dokl. Akad. Nauk SSSR **141** (1961), 1015–1018.
2. Bernstein, I. N.; Gelfand, I. M.; Ponomarev, V. A., *Coxeter functors, and Gabriel's theorem. (Russian)*, Uspehi Mat. Nauk **XXVIII 170(2)** (1973), 19–33.
3. Bovdi,V.A.; Gudivok, P.M.; Rudko, V.P., *Torsion free groups with indecomposable holonomy group I*, J. Group Theory **5** (2002), 75–96.
4. Nazarova, L. A., *Unimodular representations of the four group (Russian)*, Dokl. Akad. Nauk SSSR **140(5)** (1961), 1011–1014.
5. Nazarova, L. A., *Representations of a tetrad. (Russian)*, Izv. Akad. Nauk SSSR Ser. Mat **31** (1967), 1361–1378.

INSTITUTE OF MATHEMATICS, UNIVERSITY OF DEBRECEN
P.O. BOX 12, H-4010 DEBRECEN, HUNGARY
INSTITUTE OF MATHEMATICS AND INFORMATICS, COLLEGE OF NYÍREGYHÁZA
SÓSTÓI ÚT 31/B, H-4410 NYÍREGYHÁZA, HUNGARY
E-MAIL: VBOVDI@MATH.KLTE.HU

V.P. Rudko
DEPARTMENT OF ALGEBRA, UNIVERSITY OF UZHGOROD
88 000, UZHGOROD, UKRAINE